



Syddansk Universitet

The genetic basis for cognitive ability, memory, and depression symptomatology in middle-aged and elderly chinese twins

Xu, Chunsheng; Sun, Jianping; Ji, Fuling; Tian, Xiaocao; Duan, Haiping; Zhai, Yaoming; Wang, Shaojie; Pang, Zengchang; Zhang, Dongfeng; Zhao, Zhongtang; Li, Shuxia; Hjelmberg, Jacob v. B.; Christensen, Kaare; Tan, Qihua

Published in:

Twin Research and Human Genetics

DOI:

[10.1017/thg.2014.76](https://doi.org/10.1017/thg.2014.76)

Publication date:

2015

Document version

Publisher's PDF, also known as Version of record

Citation for pulished version (APA):

Xu, C., Sun, J., Ji, F., Tian, X., Duan, H., Zhai, Y., ... Tan, Q. (2015). The genetic basis for cognitive ability, memory, and depression symptomatology in middle-aged and elderly chinese twins. *Twin Research and Human Genetics*, 18(1), 79-85. DOI: 10.1017/thg.2014.76

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The Genetic Basis for Cognitive Ability, Memory, and Depression Symptomatology in Middle-Aged and Elderly Chinese Twins

Chunsheng Xu,^{1,2} Jianping Sun,² Fuling Ji,² Xiaocao Tian,² Haiping Duan,² Yaoming Zhai,²
Shaojie Wang,² Zengchang Pang,^{1,2} Dongfeng Zhang,³ Zhongtang Zhao,¹ Shuxia Li,⁴
Jacob v.B. Hjelmberg,^{5,6} Kaare Christensen,^{4,5,6} and Qihua Tan^{4,5,6}

¹Department of Epidemiology and Health Statistics, School of Public Health, Shandong University, Jinan, Shandong, China

²Qingdao Center for Disease Control and Prevention, Qingdao, China

³Department of Public Health, Qingdao University Medical College, Qingdao, China

⁴Department of Clinical Genetics, Odense University Hospital, Odense, Denmark

⁵Epidemiology, Biostatistics and Biodemography, Institute of Public Health, University of Southern Denmark, Odense, Denmark

⁶Danish Twin Registry and Danish Aging Research Center, Institute of Public Health, University of Southern Denmark, Odense, Denmark

The genetic influences on aging-related phenotypes, including cognition and depression, have been well confirmed in the Western populations. We performed the first twin-based analysis on cognitive performance, memory and depression status in middle-aged and elderly Chinese twins, representing the world's largest and most rapidly aging population. The sample consisted of 384 twin pairs with a median age of 50 years. Cognitive function was measured using the Montreal Cognitive Assessment (MoCA) scale; memory was assessed using the revised Wechsler Adult Intelligence scale; depression symptomatology was evaluated by the self-reported 30-item Geriatric Depression (GDS-30) scale. Both univariate and multivariate twin models were fitted to the three phenotypes with full and nested models and compared to select the best fitting models. Univariate analysis showed moderate-to-high genetic influences with heritability 0.44 for cognition and 0.56 for memory. Multivariate analysis by the reduced Cholesky model estimated significant genetic ($r_G = 0.69$) and unique environmental ($r_E = 0.25$) correlation between cognitive ability and memory. The model also estimated weak but significant inverse genetic correlation for depression with cognition (-0.31) and memory (-0.28). No significant unique environmental correlation was found for depression with other two phenotypes. In conclusion, there can be a common genetic architecture for cognitive ability and memory that weakly correlates with depression symptomatology, but in the opposite direction.

■ **Keywords:** cognition, memory, depression symptomatology, multivariate, twin model

Cognition, memory, and depression symptomatology are important determinants to maintain quality of life and independence in older people. In the literature, moderate-to-high heritability has been estimated for cognitive skill, memory, and depression symptomatology by multiple twin studies in Western countries (Finkel et al., 1995; Johansson et al., 1999; Lee et al., 2010; 2012; McGue & Christensen, 2001; 2002; Sachdev et al., 2009; Skytthe et al., 2006; Tams et al., 2009), confirming the genetic regulation of aging-related phenotypes. As the three phenotypes represent important domains of aging-associated impairment, it would be interesting to study the genetic mechanisms in the regulation of these phenotypes. Results from such analysis could

help with our understanding of the biological mechanism of mental aging and provide useful information for the

RECEIVED 17 August 2014; ACCEPTED 23 October 2014. First published online 14 January 2015.

ADDRESS FOR CORRESPONDENCE: Prof. Zengchang Pang and Zhongtang Zhao, Department of Epidemiology and Health Statistics, School of Public Health, Shandong University, Jinan, Shandong, China. E-mail: cdcpang@126.com and ztzhao@sdu.edu.cn Prof. Dongfeng Zhang, Department of Public Health, Qingdao University Medical College, Qingdao, China. E-mail: zhangdf1961@126.com

development of efficient strategies for promoting mental health and healthy aging.

In the literature of twin studies, both univariate and multivariate analyses (Rijsdijk & Sham, 2002) have been applied to a variety of disease phenotypes or traits. Different from univariate twin models, multivariate twin models explore the common genetic background shared by multiple phenotypes to infer pleiotropic genetic effects underlying the covariance of multiple phenotypes. For example, based on different assumptions, the common factor common pathway and the common factor independent pathway models have been applied to infer if there are common or independent genetic mechanisms in the regulation of metabolic phenotypes (Pang et al., 2010), behavior (Dawood et al., 2005) or psychological and psychiatric traits (Kendler et al., 2012). In aging studies, twin-based multivariate modeling has been conducted on different aging-related phenotypes using bivariate or multivariate analysis, but predominantly in Western populations (Finkel et al., 2000; Kremen et al., 2007; Ogata et al., 2014; Robitaille et al., 2012; Svedberg et al., 2009; Tucker-Drob et al., 2014); for example, the bivariate study of self-rated health and cognitive abilities (Svedberg et al., 2009), the multivariate genetic analysis of adult memory (Finkel et al., 1995), and the common genetic influence on hand strength, processing speed, and working memory (Ogata et al., 2014). Interestingly, their results, based on Western populations, indicated that common genetic factors may influence memory and processing speed and that memory, processing speed and, visuospatial ability could represent different aspects of cognitive function.

The Chinese population is the world's largest and most rapidly aging population. The aging Chinese population is potentially creating a growing burden to a society under economic transition. Studying and promoting healthy aging in the Chinese population may hold the key to improving public health in China. This is especially important considering that the current findings from Western populations may not apply to the Chinese population, who live in a different social and physical environment. This article reports results from the first twin-based multivariate modeling of cognition, memory, and depression symptomatology as part of a comprehensive investigation of genetic and environmental regulation on aging-related phenotypes in middle-aged and older Chinese twins. Results from different multivariate models will be presented, with the aim of testing our hypothesis of a common genetic and environmental background for the three phenotypes. The article ends with discussion on findings from our analysis, together with their implications.

Methods

Study Participants

The study was based at the Qingdao Twin Registry, established in 1998 at the Qingdao Center for Disease Con-

trol and Prevention (Qingdao CDC; Duan et al., 2013), the first population-based twin registry in China (Li et al., 2006; 2013). Different from Western countries with well-established registration systems, recruitment of Chinese twins is through multiple channels, especially given the lower twin birth rate (0.76%; Gan & Zheng, 2002). In this study, twins were recruited through the network of the Qingdao CDC in residential communities of the Qingdao municipality, using the residence registry, medical records, and media announcements. Complete pairs of middle-aged and older twins were recruited in 2012 and 2013. Twins who were unconscious, unable or unwilling to cooperate were excluded from sampling. The final sample consisted of 384 pairs, including 244 monozygotic (MZ, 116 male pairs, 128 female pairs) and 140 dizygotic (DZ, 42 male pairs, 39 female pairs, 59 opposite sex pairs) twin pairs aged from 33 to 80 years with a median age of 50. Altogether, there were 375 male twins with a mean age of 52.3 years and 393 female twins with a mean age of 50.9 years. Venous blood was drawn for zygosity determination for like-sex twin pairs using 16 multiple short tandem sequence repeat DNA markers (Becker et al., 1997; Tomsey et al., 2001) at the central laboratory of the Qingdao Blood Center, with correct zygosity assignment ascertained to be 99.9%.

The study was approved by the Regional Ethics Committee of the Qingdao CDC Institutional Review Boards and conducted according to the principles of the Helsinki Declaration. All participants signed a consent form and completed a questionnaire and health examination at the local service center of the Qingdao CDC or at community hospital/clinics.

Assessment of Cognition, Memory, and Depression Symptomatology

For each twin pair, face-to-face interviews with each twin were performed by the same well-trained and experienced interviewer. We used the MoCA (www.mocatest.org) scale to test the general cognitive performance of the participants. The MoCA is the most age-sensitive and effective test for assessing cognitive ability of healthy aging (Gluhm et al., 2013; Ismail et al., 2010; 2013; Luis et al., 2009; Nasreddine et al., 2005). The MoCA is a brief 0–30 point assessment of global cognition that evaluates a broad array of cognitive domains including attention, orientation, abstraction, linguistic skill, delayed recall and executive ability. A recent study reported that premorbid IQ influences MoCA in both healthy subjects and patients with cognitive impairments (Alves et al., 2013). Here, we used the MoCA adjusted for education, with educational level quantified by totaling the number of years to complete the participant's highest of schooling, adding 1 score if it was less than 12 years of education. A higher cognitive score meant better cognition.

Memory was measured using the forward and backward digit span task from the Wechsler Adult Intelligence scale — Revised for China (WAIS-RC; Gong, 1989). In the

TABLE 1
Descriptive Statistics, Regression Outputs and ICCs for the Three Phenotypes

Phenotypes	Median	Quantile (%) 2.5–97.5	Regression coefficients						ICCs		
			Age	p-value	Gender	p-value	Age* Gender	p-value	MZ (95%CI)	DZ (95%CI)	p-value
Cognition	22	8.2–28.0	-4.237	5.0e-4	-2.181	0.959	0.345	0.674	0.55(0.44,0.64)	0.35(0.19,0.49)	0.018
Memory	12	6–16	-0.219	8.93e-9	-3.627	0.007	0.084	0.001	0.49(0.38,0.59)	0.29(0.12,0.44)	0.027
Depression symptomatology	7	1–22	0.036	.687	1.460	0.642	-0.027	0.657	0.45(0.34,0.55)	0.35(0.19,0.49)	0.265

Note: Bold type indicates statistical significance ($p = 0.05$).

forward digit condition, the interviewer read aloud a series of numbers of increasing length, and the participant was instructed to repeat the numbers in the same order. For the backward digits, the interviewer also read aloud a series of numbers of increasing length, but the participant was instructed to repeat the numbers in the reverse order. The sequences were increased in length by 1 unit in each subsequent trial until the participant failed two trials in a row of the same sequence length. The forward digit battery consisted of sequences of 3–9 units, and the backward digits test consisted of sequences of 2–8 units. Forward and backward scores were based on the longest length of correct answers in each condition. Memory was measured as the sum of forward and backward digit span scores (range, 0–17). A higher digit span score indicated better memory performance.

Depression symptomatology was assessed by the self-reported GDS-30 (Brink et al., 1982). All the items were transformed so that the higher the total score, the more severe the participant's mental condition.

Statistical Analysis

Data were entered and corrected by Epidata3.1 (www.epidata.dk). Considering the skewed distributions of cognition, memory, and depression symptomatology, all three measurements were transformed by Box–Cox power transformation (Box & Cox, 1964) to ensure approximately normal distribution using the free R package *car* (<http://cran.r-project.org/web/packages/car/index.html>). Data were analyzed to estimate the intra-pair correlation coefficients (ICCs) and test statistical significance for the effects of age and sex using *twinlm* function in the free R package *met*s (<http://cran.r-project.org/web/packages/mets/index.html>).

The classical twin models were applied to each of the three phenotypes to decompose the observed variance into the additive genetic (A), shared environmental (C), and unique environmental (E) components by fitting models encompassing the A, C, and E components (the full ACE model) or the A, D, and E components (the full ADE model) with the ADE model only considered when phenotype correlation in MZ twins was over two times that in the DZ twins. By dropping each variance component from the full model in turn, the nested models (AE, CE, DE, E) were fitted and statistical significance of the corresponding removed

variance component was tested using the likelihood ratio test. This was followed by the multivariate analysis (Rijsdijk & Sham, 2002) that fitted a full multivariate Cholesky model to all phenotypes and estimated genetic correlation (r_G), common environmental correlation (r_C) and unique environmental correlation (r_E) among the three phenotypes. As a balance between model fit and parsimony in the number of variables, Akaike's Information Criterion (AIC; Akaike, 1987) was used to determine the suitability of models, with the lowest value indicating the best model.

In all model fitting, age and sex (1 for males and 2 for females) were included as the covariates to adjust for their effects on the three phenotypes. Robustness of analysis was assessed using bootstrap resampling to calculate empirical 95% CIs for estimated variables. All the multivariate twin modeling was performed by the free software Mx package (<http://www.vcu.edu/mx>).

Results

Descriptive Statistics

The descriptive statistics (median, percentiles) for the three phenotypes are shown in Table 1, together with results from fitting linear models to each measurement for the effects of age and sex. There was a strong pattern of decline in cognition and memory with increasing age, as indicated by the highly significant negative regression coefficients. For sex, memory was significantly lower in females than in males but there was a significant age–sex interaction in favor of females, meaning that with increasing age, females gradually outperform males in memory. Age and sex had no significant influence on depression symptomatology ($p > 0.05$). Table 1 also displays the estimated ICCs for MZ and DZ twins separately. For cognition and memory, the ICC was significantly higher for MZ than for DZ twins, suggesting genetic influences. The ICC for depression symptomatology was also higher in MZ than in DZ twin pairs but the difference failed to reach statistical significance.

Univariate Twin Models

In Table 1, the estimated ICCs for MZ twins were all lower than the doubled ICCs for DZ twins, thus no ADE model was considered in model fitting. Heritability first estimated by the univariate ACE models was low (0.22 for cognition, 95% CI: -0.13–0.57; 0.20 for depression symptomatology,

TABLE 2**The Full and the Best Fitting Model for Cognition, Memory, and Depression Symptomatology Adjusted for Age and Gender**


Phenotype	Model	A (95%CI)	C (95%CI)	E (95%CI)	-2LL	AIC	χ^2	Δdf	p value
Full model									
Cognition	ACE	0.2,206(-0.1,306,0.5,718)	0.2,009(-0.1,102,0.5,120)	0.5,785(0.4,786,0.6,783)	8,578.376	8,592.377	—	—	—
Memory	ACE	0.4,767(0.1,592,0.7,941)	0.0812(-0.2,063,0.3,686)	0.4,422(0.3,567,0.5,277)	5,088.094	5,100.094	—	—	—
Depression symptomatology	ACE	0.2,011(-0.1,338,0.5,361)	0.2,507(-0.0477,0.5,492)	0.5,482(0.4,530,0.6,433)	4,761.194	4,773.195	—	—	—
Best fitting model									
Cognition	AE	0.4,366(0.3,436,0.5,296)	—	0.5,634(0.4,704,0.6,564)	8,579.842	8,591.842	1.466	1	0.2,260
Memory	AE	0.5,629(0.4,812,0.6,446)	—	0.4,371(0.3,554,0.5,188)	5,088.388	5,098.388	0.294	1	0.5,877
Depression symptomatology	CE	—	0.4,176(0.3,348,0.5,003)	0.5,825(0.4,997,0.6,652)	4,762.660	4,772.661	1.466	1	0.2,260

Note: A = Additive genetic influence, C = Shared environmental influence, E = Unique/non-shared environment influence, 2LL = two times the log-likelihood; Δdf = difference in degree of freedom.

95% CI: -0.13–0.54) to moderate (0.48 for memory, 95% CI: 0.16–0.79; Table 2). Model comparison between the full ACE models and their nested models selected the best fitting models as the AE model for cognition and memory and the CE model for depression symptomatology. The best fitting AE models estimated moderate-to-high heritability for cognition (0.44, 95% CI: 0.34–0.53) and memory (0.56, 95% CI: 0.48–0.64) while the best-fitting CE model for depression symptomatology estimated a strong common environmental component (0.42, 95% CI: 0.33–0.50). In contrast to the genetic component, the unique environment had a relatively high contribution to the variation of the three phenotypes in all the fitted models.

Multivariate Cholesky Models

We started multivariate analysis by fitting a full Cholesky decomposition model to the three phenotypes. In the full model, the 95% confidence intervals for the estimated common environmental contributions to total variance or covariance either overlapped with zero or were very close to zero. With this consideration, we fitted a reduced Cholesky decomposition model by fixing loadings for common environmental factors to zero. Comparison of the reduced (-2LLK = 5,811.526, df = 2,273) with the full (-2LLK = 5,818.894, df = 2,279) Cholesky model using likelihood ratio test gave a p -value of 0.29 (χ^2 = 7.37, df = 6). As a result, the reduced Cholesky model was preferred with consideration of model parsimony and limited sample size. Figure 1 is the path diagram for the reduced Cholesky model, with each factor having one loading less than the previous one. Along the path, arrows are the standardized path coefficients for the additive genetic and unique environmental factors. The estimated path coefficients show strong factor loading from the genetic factor of cognition to memory. In contrast, the unique environmental factors for cognition and memory have only minor loading to depression symptomatology. In Table 3, we show the estimated genetic and unique environmental correlation coefficients among the three phenotypes. As expected, the model estimated a high genetic correlation (r_G = 0.69, 95% CI: 0.57–0.79)

**FIGURE 1.**

(Colour online) Path diagram of the reduced Cholesky decomposition model. The standardized path coefficients show strong factor loading from the genetic factor of cognition to memory. In contrast, the unique environmental factors for cognition and memory have only minor loading to depression symptomatology.

between cognition and memory. Depression had low but significant genetic correlation with cognition (r_G = -0.31, 95% CI: -0.46– -0.14) and memory (r_G = -0.28, 95% CI: -0.44– -0.11), suggesting their inverse genetic relationship with depression. The unique environmental correlation was only significant between cognition and memory (r_E = 0.25, 95% CI: 0.14–0.36).

Discussion

We have fitted both univariate and multivariate twin models to three important domains of aging phenotypes, that is, cognition, memory, and depression symptomatology. Our best fitting model estimated a moderate heritability for cognition (0.44; Table 2). This estimate is relative lower than some of the reported studies but is comparable with results from a Danish twin study (McGue & Christensen, 2001). In a recent study, Karlsgodt et al. estimated heritability for

TABLE 3**Genetic and Environmental Correlation Coefficients for the Reduced Cholesky Decomposition Model**

Phenotype	Cognition	Memory	Depression symptomatology
A			
Cognition	1.000		
Memory	0.685 (0.570,0.788)*	1.000	
Depression symptomatology	-0.305 (-0.462,-0.143)*	-0.281 (-0.444,-0.113)*	1.000
E			
Cognition	1.000		
Memory	0.249 (0.135,0.357)*	1.000	
Depression symptomatology	-0.043 (-0.159,0.074)	-0.040 (-0.156,0.078)	1.000

Note: **p* value .05.

a subtest of the WAIS of 0.54 for forward digit span using family data (Karlsgodt et al., 2010). Interestingly, this is very close to our estimate of 0.56 by the best fitting model for memory. Different from cognition and memory, our best fitting model for depression estimated no genetic effect in depression symptomatology. Based on a large sample of Danish twins, Johnson et al. (2002) studied the additive genetic effects on depression symptoms with moderate heritability estimates (affective scale $h^2 = 0.27$, somatic scale $h^2 = 0.26$, total $h^2 = 0.29$). Their estimates are, however, close to our estimate of 0.20 (95% CI: -0.13–0.54) although the latter is statistically insignificant. We think that the discrepancy in the estimated genetic effect in depression symptomatology could be due to our limited power in estimating low heritability using a small twin sample, rather than population differences.

The focus of our multivariate analysis is to examine the common genetic and environmental background of the three phenotypes. Our estimates revealed a strong genetic correlation between cognition and memory, suggesting common genetic basis of the two phenotypes. In our assessment instrument, MoCA-based cognition is a composite measurement that evaluates a broad array of cognitive domains, including attention, orientation, abstraction, linguistic skill, delayed recall, and executive ability. The WAIS-RC (Finkel et al., 1995; Johansson et al., 1999) measures memory with a backward and forward digit-span to assess an individual's intelligence in retrieving information. The WAIS backward and forward digit span test measures attention and short-term memory while the MoCA test includes items for delayed recall and attention. Thus, it is not surprising to see from our results that the two domains are highly genetically similar. Using results on twins, Finkel and McGue (1993) reported that the relationship between cognition and memory was genetic in nature, a conclusion that is consistent with our finding from multivariate analysis. Overall, we conclude that memory and cognitive ability are highly genetically interrelated and if one becomes impaired, the other could also be affected.

As can be seen from Table 3, depression symptomatology has a weak but significant genetic correlation with cognition and memory. The negative and low genetic correlation coefficients imply that some genes that affect cognition and

memory could also affect depression symptom but in an opposite direction. Although the GDS-30 is an assessment instrument to identify depression in the older people, different from cognition and memory, it does not seem to change by age in the age range of our samples (Table 1). Considering that the median age of our samples is 50 years, the reported high peak of depression risk in older subjects (80 years or older; Mirowsky & Ross, 1992) was actually missed. It would be interesting to study older aged Chinese twins, which could perhaps reveal a different picture of the multivariate models.

Indeed, our study was limited by its small sample size and relatively younger age. However, even with a relatively limited and middle-to-older aged samples, our study found interesting genetically correlated and uncorrelated patterns among the three important domains of aging-associated phenotypes, although with considerably large confidence intervals. The results have encouraged us to extend our study to older aged twins with the hope of capturing the dynamic genetic and environmental regulation patterns in mental aging. As shown in Table 1, both cognition and memory have a strong declining pattern with age in our middle-to-older aged twins. As a continuation of our current project, we are going to set up a longitudinal investigation to measure changes in aging phenotypes. The longitudinal study could provide important information concerning the genetic and environmental contribution to the rate of change over time, on top of the mean phenotype levels, and help to explain why and how people age differently.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (grant number 31371024). All authors contributed significantly to the study with Chunsheng Xu, Zengchang Pang, Dongfeng Zhang, Qihua Tan: experiment design, conduct of project, data analysis, drafting of manuscript; Jianping Sun, Fuling Ji, Xiaocao Tian, Haiping Duan, Yaoming Zhai, Shaojie Wang, Zhongtang Zhao: conduct of project, sample collection, data preparation; Shuxia Li, Jacob v.B. Hjelmberg, Kaare Christensen: analysis plan, interpretation, manuscript revision. We also thank the twins for their volunteer participation in this study.

References

- Akaike, H. (1987). Factor analysis' and AIC. *Psychometrika*, 52, 317–332.
- Alves, L., Simoes, M. R., Martins, C., Freitas, S., & Santana, I. (2013). Premorbid IQ influence on screening tests' scores in healthy patients and patients with cognitive impairment. *Journal of Geriatric Psychiatry and Neurology*, 26, 117–126.
- Becker, A., Busjahn, A., Faulhaber, H. D., Bahring, S., Robertson, J., Schuster, H., . . . Luft, F. C. (1997). Twin zygosity: Automated determination with microsatellites. *Journal of Reproductive Medicine*, 42, 260–266.
- Box, G. E. P., & Cox, D. R. (1964). An analysis of transformations. *Journal of the Royal Statistical Society, Series B*, 26, 211–252.
- Brink, T. A., Yesavage, J. A., Lum, O., Heersema, P., Adey, M., & Rose, T. L. (1982). Screening tests for geriatric depression. *Clinical Gerontologist*, 1, 37–44.
- Dawood, K., Kirk, K. M., Bailey, J. M., Andrews, P. W., & Martin, N. G. (2005). Genetic and environmental influences on the frequency of orgasm in women. *Twin Research and Human Genetics*, 8, 27–33.
- Duan, H., Ning, F., Zhang, D., Wang, S., Zhang, D., Tan, Q., . . . Pang, Z. (2013). The qingdao twin registry: A status update. *Twin Research and Human Genetics*, 16, 79–85.
- Finkel, D., & McGue, M. (1993). The origins of individual differences in memory among the elderly: A behavior genetic analysis. *Psychology and Aging*, 8, 527–537.
- Finkel, D., Pedersen, N., & McGue, M. (1995). Genetic influences on memory performance in adulthood: comparison of Minnesota and Swedish twin data. *Psychology and Aging*, 10, 437–446.
- Finkel, D., Pedersen, N. L., & Harris, J. R. (2000). Genetic mediation of the associations among motor and perceptual speed and adult cognitive abilities. *Aging, Neuropsychology, and Cognition (Neuropsychology, Development and Cognition: Section B)*, 7, 141–155.
- Gan, J., & Zheng, J. (2002). Regional distribution of birth rate of twins in China. *China Public Health*, 18, 658–659.
- Gluhm, S., Goldstein, J., Loc, K., Colt, A., Liew, C. V., & Corey-Bloom, J. (2013). Cognitive performance on the minimal state examination and the montreal cognitive assessment across the healthy adult lifespan. *Cognitive and Behavioral Neurology*, 26, 1–5.
- Gong, X. Y. (1989). *Manual for the Wechsler Adult Intelligence Scale: Revised in China (WAIS-RC)*. Changsha, Hunan, China: Hunan Medical University.
- Ismail, Z., Mulsant, B. H., Herrmann, N., Rapoport, M., Nilsson, M., & Shulman, K. (2013). Canadian academy of geriatric psychiatry survey of brief cognitive screening instruments. *Canadian Geriatrics Journal*, 16, 54–60.
- Ismail, Z., Rajji, T. K., & Shulman, K. I. (2010). Brief cognitive screening instruments: An update. *International Journal of Geriatric Psychiatry*, 25, 111–120.
- Johansson, B., Whitfield, K., Pedersen, N. L., Hofer, S. M., Ahern, F., & McClearn, G. E. (1999). Origins of individual differences in episodic memory in the oldest-old: A population-based study of identical and same-sex fraternal twins aged 80 and older. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 54, P173–P179.
- Johnson, W., McGue, M., Gaist, D., Vaupel, J. W., & Christensen, K. (2002). Frequency and heritability of depression symptomatology in the second half of life: Evidence from Danish twins over 45. *Psychological Medicine*, 32, 1175–1185.
- Karlsodt, K. H., Kochunov, P., Winkler, A. M., Laird, A. R., Almasy, L., Duggirala, R., . . . Glahn, D. C. (2010). A multimodal assessment of the genetic control over working memory. *The Journal of Neuroscience*, 30, 8197–8202.
- Kendler, K. S., Aggen, S. H., & Patrick, C. J. (2012). A multivariate twin study of the DSM-IV criteria for antisocial personality disorder. *Biological Psychiatry*, 71, 247–253.
- Kremen, W. S., Jacobsen, K. C., Xian, H., Eisen, S. A., Eaves, L. J., Tsuang, M. T., . . . Lyons, M. J. (2007). Genetics of verbal working memory processes: A twin study of middle-aged men. *Neuropsychology*, 21, 569–580.
- Lee, T., Henry, J. D., Trollor, J. N., & Sachdev, P. S. (2010). Genetic influences on cognitive functions in the elderly: A selective review of twin studies. *Brain Research Reviews*, 64, 1–13.
- Lee, T., Mosing, M. A., Henry, J. D., Trollor, J. N., Ames, D., Martin, N. G., . . . Oats Research. (2012). Genetic influences on four measures of executive functions and their covariation with general cognitive ability: The older Australian twins study. *Behavior Genetics*, 42, 528–538.
- Li, L., Gao, W., Lv, J., Cao, W., Zhan, S., Yang, H., . . . Hu, Y. (2006). Current status of the Chinese national twin registry. *Twin Research and Human Genetics*, 9, 747–752.
- Li, L., Gao, W., Yu, C., Lv, J., Cao, W., Zhan, S., . . . Hu, Y. (2013). The Chinese national twin registry: An update. *Twin Research and Human Genetics*, 16, 86–90.
- Luis, C. A., Keegan, A. P., & Mullan, M. (2009). Cross validation of the montreal cognitive assessment in community dwelling older adults residing in the Southeastern US. *International Journal of Geriatric Psychiatry*, 24, 197–201.
- McGue, M., & Christensen, K. (2001). The heritability of cognitive functioning in very old adults: Evidence from Danish twins aged 75 years and older. *Psychology and Aging*, 16, 272–280.
- McGue, M., & Christensen, K. (2002). The heritability of level and rate-of-change in cognitive functioning in Danish twins aged 70 years and older. *Experimental Aging Research*, 28, 435–451.
- Mirowsky, J., & Ross, C. E. (1992). Age and depression. *Journal of Health and Social Behavior*, 33, 187–205.
- Nasreddine, Z. S., Phillips, N. A., Bedirian, V., Charbonneau, S., Whitehead, V., Collin, I., . . . Chertkow, H. (2005). The montreal cognitive assessment, MoCA: A brief screening tool for mild cognitive impairment. *Journal of the American Geriatrics Society*, 53, 695–699.
- Ogata, S., Kato, K., Honda, C., & Hayakawa, K. (2014). Common genetic factors influence hand strength, processing speed, and working memory. *Journal of Epidemiology*, 24, 31–38.

- Pang, Z., Zhang, D., Li, S., Duan, H., Hjelmberg, J., Kruse, T. A., . . . Tan, Q. (2010). Multivariate modelling of endophenotypes associated with the metabolic syndrome in Chinese twins. *Diabetologia*, 53, 2554–2561.
- Rijsdijk, F. V., & Sham, P. C. (2002). Analytic approaches to twin data using structural equation models. *Briefings in Bioinformatics*, 3, 119–133.
- Robitaille, A., Muniz, G., Piccinin, A. M., Johansson, B., & Hofer, S. M. (2012). Multivariate longitudinal modeling of cognitive aging: Associations among change and variation in processing speed and visuospatial ability. *Journal of Gerontopsychology and Geriatric Psychiatry*, 25, 15–24.
- Sachdev, P. S., Lammel, A., Trollor, J. N., Lee, T., Wright, M. J., Ames, D., . . . Oats Research Team. (2009). A comprehensive neuropsychiatric study of elderly twins: The older Australian twins study. *Twin Research and Human Genetics*, 12, 573–582.
- Skytthe, A., Kyvik, K., Bathum, L., Holm, N., Vaupel, J. W., & Christensen, K. (2006). The Danish twin registry in the new millennium. *Twin Research and Human Genetics*, 9, 763–771.
- Svedberg, P., Gatz, M., & Pedersen, N. L. (2009). Genetic and environmental mediation of the associations between self-rated health and cognitive abilities. *Experimental Aging Research*, 35, 178–201.
- Tambs, K., Ronning, T., Prescott, C. A., Kendler, K. S., Reichborn-Kjennerud, T., Torgersen, S., . . . Harris, J. R. (2009). The norwegian institute of public health twin study of mental health: Examining recruitment and attrition bias. *Twin Research and Human Genetics*, 12, 158–168.
- Tomsey, C. S., Kurtz, M., Kist, F., Hockensmith, M., & Call, P. (2001). Comparison of PowerPlex 16, PowerPlex1.1/2.1, and ABI AmpflSTR Profiler Plus/COfiler for forensic use. *Croatian Medical Journal*, 42, 239–243.
- Tucker-Drob, E. M., Reynolds, C. A., Finkel, D., & Pedersen, N. L. (2014). Shared and unique genetic and environmental influences on aging-related changes in multiple cognitive abilities. *Developmental Psychology*, 50, 152–166.